Index of Authors

ABITHA DEVI, N. *see* VEERASWAMY, M. 807–809

ABNEY, E. R., EVANS, W. H. & PARKHOUSE, R. M. E. Location of nucleotide pyrophosphatase and alkaline phosphodiesterase activities on the lymphocyte surface membrane 293–299

ADAMS, P. *see* HARDINGHAM, T. E. 143–147

ADAMS, P. A. The kinetics and mechanism of the recombination reaction between apomyoglobin and haemin 371–376

ADLER, L. S. *see* GOLD, R. J. M. 157–160

AKHTAR, M. *see* PEARCE, C. J. 601–606

ALLEN, J. D. & THOMA, J. A. Subsite mapping of enzymes. Application of the depolymerase computer model to two α-amylases 121–131

ALLEN, J. D. & THOMA, J. A. Subsite mapping of enzymes. Depolymerase computer modelling 105–120

AMBLER, R. P. *see* VAN BEEUMEN, J. 757–769

ANASTASI, A. *see* BANNISTER, J. V. 35–42

ANDERSON, P. J. Chicken muscle and fibroblast actin structure 185–187

ANTHONY, C. *see* PEARCE, C. J. 601–606

APPS, D. K. & GLEED, C. D. Interaction of pigeon liver nicotinamide–adenine dinucleotide kinase with Cibacron Blue F3GA 441–443

ARMSTRONG, J. MCD. & TRENTHAM, D. R. The reactions of D-glyceraldehyde 3-phosphate with thiols and the holoenzyme of D-glyceraldehyde 3-phosphate dehydrogenase and of inorganic phosphate with the acyl-holoenzyme 513–527

ATASSI, M. Z. *see* LEE, C.-L. 89–93

ATFIELD, G. *see* HOLNESS, N. J. 15–22

BANIK, N. L., GOHIL, K. & DAVISON, A. N. The action of snake venom, phospholipase A and trypsin on purified myelin *in vitro* 273–277

BANNISTER, J. V., BANNISTER, W. H., ANASTASI, A. & WOOD, E. J. Isolation, characterization and oxygen equilibrium of an extracellular haemoglobin from *Eunice aphroditois* (Pallas) 35–42

BANNISTER, W. H. *see* BANNISTER, J. V. 35–42

BARNETT, J. E. G. *see* PEARCE, C. J. 601–606

BARON, A. J. *see* WOOTTON, J. C. 803–806

BEELEY, J. G. Location of the carbohydrate groups of ovomucoid 335–345

BENVENISTE, P. *see* BOUVIER, P. 267–271

BERNINGER, R. W. & MATHIS, R. K. Isolation and characterization of α-1-antitrypsin from rhesus-monkey serum 95–104

BETTS, S. A. *see* WALKER, J. H. 355–362

BEUTLER, E., YOSHIDA, A., KUHL, W. & LEE, J. E. S. The subunits of human hexosaminidase A 541–543

BLOXHAM, D. P. & CHALKLEY, R. A. Selective modification of rabbit muscle pyruvate kinase by 5-chloro-4-oxopentanoic acid 201–211

BLOXHAM, D. P. *see also* CHALKLEY, R. A. 213–219

BOLANDER, F. F., JR. & FELLOWS, R. E. The purification and characterization of rabbit placental lactogen 775–782

BOOTH, A. G. & KENNY, A. J. Proteins of the kidney microvillus membrane. Identification of subunits after sodium dodecyl sulphate/polyacrylamide-gel electrophoresis 395–407

BOTTOMLEY, R. C., STORER, A. C. & TRAYER, I. P. A new method of quantitative affinity chromatography and its application to the study of myosin 667–676

BOUILLON, R., VAN BAELEN, H. & DE MOOR, P. The transport of vitamin D in the serum of primates 463–466

BOUVIER, P., ROHMER, M., BENVENISTE, P. & OURISSON, G. $\Delta^{8(14)}$-Steroids in the bacterium *Methylococcus capsulatus* 267–271

BRITTAIN, T. & GREENWOOD, C. Kinetic studies on [methionine sulphoxide]cytochrome *c* 621–626

BRITTAIN, T., SPRINGALL, J., GREENWOOD, C. & THOMSON, A. J. Low-temperature studies on mixed-valence cytochrome oxidase by using magnetic circular dichroism. Evidence for haem–haem interactions 811–813

BROCKLEHURST, K. & CORNISH-BOWDEN, A. The pre-eminence of $k_{cat.}$ in the manifestation of optimal enzymic activity delineated by using the Briggs–Haldane two-step irreversible kinetic model 165–166

BROCKLEHURST, K. *see also* MALTHOUSE, J. P. G. 221–234; NORRIS, R. 245–257; SHIPTON, M. 235–244

BRODSKY-DOYLE, B., LEONARD, K. R. & REID, K. B. M. Circular-dichroism and electron-microscopy studies of human subcomponent C1q before and after limited proteolysis by pepsin 279–286

BROWN, J. S. *see* MURRAY, K. 317–322

BROWN, S. B. Stereospecific haem cleavage. A model for the formation of bile-pigment isomers *in vivo* and *in vitro* 23–27

BRUCE, S. A. *see* MURRAY, K. 317–322

CAMPBELL, A. M. The effects of deoxyribonucleic acid secondary structure on tertiary structure 615–620

CHALKLEY, R. A. & BLOXHAM, D. P. Identification of cysteine as the reactive group in pyruvate kinase alkylated by 5-chloro-4-oxopentanoic acid 213–219

CHALKLEY, R. A. *see also* BLOXHAM, D. P. 201–211

CHAPMAN, K. S. R. & INGLE, J. The stability, polyadenylic acid content and ribonucleoprotein form of nuclear ribonucleic acid in artichoke 585–600

CHENG, K.-W. Carboxymethylation of methionine residues in bovine pituitary luteinizing hormone and its subunits. Effects on the binding activity with receptor sites and interactions between subunits 71–77

CHENG, K.-W. Carboxymethylation of methionine residues in bovine pituitary luteinizing hormone and its subunits. Location of specifically modified methionine residues 79–87

CHENG, K.-W. Purification and properties of bovine pituitary follitropin 651–659

CHIERICATO, G. *see* GIGLIO, J. R. 29–33

CHRISTELLER, J. T. *see* LAING, W. A. 563–570

CHRISTIE, W. W., HUNTER, M. L. & VERNON, R. G. Triacylglycerol biosynthesis in rat adipose-tissue homogenates 571–577

CLARKE, F. M. & MORTON, D. J. Aldolase binding to actin-containing filaments. Formation of paracrystals 797–798

CLEMMENSEN, I. *see* MÜLLERTZ, S. 545–553

CORBIN, J. D. *see* SUGDEN, P. H. 409–422, 423–437

CORNISH-BOWDEN, A. Algebraic methods for deriving steady-state rate equations. Practical difficulties with mechanisms that contain repeated rate constants 167
CORNISH-BOWDEN, A. *see also* BROCKLEHURST, K. 165–166; STORER, A. C. 1–5, 7–14
COX, R. A. *see* HIRST, W. 259–265

DANSON, M. J. & PERHAM, R. N. Evidence for two lipoic acid residues per lipoate acetyltransferase chain in the pyruvate dehydrogenase multienzyme complex of *Escherichia coli* 677–681
DAVIDSON, I. W., SUTHERLAND, I. W. & LAWSON, C. J. Purification and properties of an alginate lyase from a marine bacterium 707–713
DAVISON, A. N. *see* BANIK, N. L. 273–277
DE LUCA, L. M. *see* FROT-COUTAZ, J. P. 799–801
DE MOOR, P. *see* BOUILLON, R. 463–466
DELLA PIETRA, G. *see* PORTA, R. 579–583
DIXON, H. B. F. Removal of a bound ligand from a macromolecule by gel filtration 161–162
DREWE, R. H. & WINZOR, D. J. An electrophoretic study of the reversible binding of phosphate to ovalbumin 737–741
DUANCE, V. C. *see* YATES, D. W. 719–728

EGAMI, F. *see* HATANAKA, H. 445–448
ELLIOTT, A. The instantaneous monitoring of polyacrylamide gels during electrophoresis 743–748
ENGLBERGER, F. M. *see* OLSON, J. M. 769–774
ESPOSITO, C. *see* PORTA, R. 579–583
EVANS, W. H. *see* ABNEY, E. R. 293–299

FELDHOFF, R. C. & PETERS, T., JR. Determination of the number and relative position of tryptophan residues in various albumins 529–533
FELLOWS, R. E. *see* BOLANDER, F. F., JR. 775–782
FERGUSON, S. J., LLOYD, W. J. & RADDA, G. K. A method for determining the adenosine triphosphatase content of energy-transducing membranes. Reaction of 4-chloro-7-nitrobenzofurazan with the adenosine triphosphatase of bovine heart submitochondrial particles 347–353
FITCH, W. M. *see* TARR, G. E. 193–199
FOTTRELL, P. F. *see* PIGGOTT, C. O. 715–717
FREEDMAN, R. B. *see* HAWKINS, H. C. 385–393; IBBETSON, A. L. 377–384
FROT-COUTAZ, J. P. & DE LUCA, L. M. Chemical synthesis of all-*trans*-β-retinoyl phosphate 799–801

GAUTHIER, F. & MOURAY, H. Rat α_2 acute-phase macroglobulin. Isolation and physicochemical properties 661–665
GERO, S. D. *see* PEARCE, C. J. 601–606
GIGLIO, J. R., ROSSI, A., LEONE, F. A., CHIERICATO, G. & SAY, J. C. Isolation and characterization of an active three-chain molecular species of bovine thrombin 29–33
GILMAN, J. G. Mouse haemoglobin beta chains. Comparative sequence data on adult major and minor beta chains from two species, *Mus musculus* and *Mus cervicolor* 43–53
GLEED, C. D. *see* APPS, D. K. 441–443
GOHIL, K. *see* BANIK, N. L. 273–277
GOLD, R. J. M., TENENHOUSE, H. S. & ADLER, L. S. A method for calculating the relative protein contents of the major keratin components from their amino acid composition 157–160
GOLD, R. J. M. *see also* TENENHOUSE, H. S. 149–157
GORDON, A. H. The α macroglobulins of rat serum 643–650
GRAND, R. J. A., WILKINSON, J. M. & MOLE, L. E. The amino acid sequence of rabbit cardiac troponin I 633–641
GREEN, N. M. *see* KNIGHT, C. G. 323–333
GREENWOOD, C. *see* BRITTAIN, T. 621–626; 811–813
GRUNWALD, M. & HILL, H. Z. Characterization of the glucose 6-phosphate dehydrogenase activity in rat liver mitochondria 683–687

HALL, D. O. *see* HASE, T. 55–63
HARDINGHAM, T. E. & ADAMS, P. A method for the determination of hyaluronate in the presence of other glycosaminoglycans and its application to human intervertebral disc 143–147
HASE, T., OHMIYA, N., MATSUBARA, H., MULLINGER, R. N., RAO, K. K. & HALL, D. O. Amino acid sequence of a four-iron–four-sulphur ferredoxin isolated from *Bacillus stearothermophilus* 55–63
HATANAKA, H., OGAWA, Y. & EGAMI, F. Arylsulphatase and glycosulphatase of *Charonia lampas*. Substrate specificity towards sugar sulphate derivatives 445–448
HAWKINS, H. C. & FREEDMAN, R. B. Thiol–protein disulphide oxidoreductases. Differences between protein disulphide-isomerase and glutathione–insulin transhydrogenase activities in ox liver 385–393
HAYWARD, C. F. *see* OFFORD, R. E. 480–486
HERRING, G. M. A comparison of bone matrix and tendon with particular reference to glycoprotein content 749–755
HICK, P. E. *see* TURNER, A. J. 819–822
HILL, H. Z. *see* GRUNWALD, M. 683–687
HIRST, W. & COX, R. A. The construction and analysis of sucrose gradients for use with zonal rotors 259–265
HODSON, A. W. *see* LATNER, A. L. 697–705
HOLLADAY, L. A. *see* SUGDEN, P. H. 409–422
HOLNESS, N. J. & ATFIELD, G. Application of a spectrophotometric method to the determination of the composition of oligonucleotides obtained from cysteine transfer ribonucleic acid 15–22
HUGHES, S. G. *see* MURRAY, K. 317–322
HUNTER, M. L. *see* CHRISTIE, W. W. 571–577

IBBETSON, A. L. & FREEDMAN, R. B. Thiol–protein disulphide oxidoreductases. Assay of microsomal membrane-bound glutathione–insulin transhydrogenase and comparison with protein disulphide-isomerase 377–384
INGLE, J. *see* CHAPMAN, K. S. R. 585–600

JENSEN, J. & OTTOLENGHI, P. Adenosine diphosphate binding to sodium-plus-potassium ion-dependent adenosine triphosphatase. The role of lipid in the nucleotide–potassium ion interplay 815–817
JOHNSON, W. H. *see* OFFORD, R. E. 480–486
JONES, D. S. *see* OLD, J. M. 503–511

KAMEN, M. D. *see* VAN BEEUMEN, J. 757–769
KAPLAN, H. *see* MALCHY, B. L. 173–175
KAVANAGH, J. P. *see* WOOTTON, J. C. 803–806
KENNY, A. J. *see* BOOTH, A. G. 395–407
KIDO, R. *see* NOGUCHI, T. 607–613
KNIGHT, C. G. & GREEN, N. M. The accessibility of protein-bound dinitrophenyl groups to univalent fragments of anti-dinitrophenyl antibody 323–333
KRISHNAN KUTTY, R. *see* VEERASWAMY, M. 807–809
KUHL, W. *see* BEUTLER, E. 541–543

LAING, W. A. & CHRISTELLER, J. T. A model for the kinetics of activation and catalysis of ribulose 1,5-bisphosphate carboxylase 563–570
LATNER, A. L. & HODSON, A. W. Human liver alkaline phosphatase purified by affinity chromatography, ultracentrifugation and polyacrylamide-gel electrophoresis 697–705
LAWSON, C. J. *see* DAVIDSON, I. W. 707–713
LEDERER, B. & STALMANS, W. Human liver glycogen phosphorylase. Kinetic properties and assay in biopsy specimens 689–695
LEE, C.-L. & ATASSI, M. Z. Delineation of the third antigenic site of lysozyme by application of a novel 'surface-simulation' synthetic approach directly linking the conformationally adjacent residues forming the site 89–93
LEE, J. E. S. & YOSHIDA, A. Purification and chemical characterization of human hexosaminidases A and B 535–539
LEE, J. E. S. *see also* BEUTLER, E. 541–543
LEONARD, K. R. *see* BRODSKY-DOYLE, B. 279–286
LEONE, F. A. *see* GIGLIO, J. R. 29–33
LICHTENSTEIN, A. V. & SHAPOT, V. S. A model for cytoplasm-governed regulation 783–794
LLOYD, W. J. *see* FERGUSON, S. J. 347–353
LONDESBOROUGH, J. *see* VARIMO, K. 363–370
LOVETT, M. G. *see* WOOTTON, J. C. 803–806

MALCHY, B. L. & KAPLAN, H. Unmasking of histone amino groups in chromatin at high pH 173–175
MALTHOUSE, J. P. G. & BROCKLEHURST, K. Preparation of fully active ficin from *Ficus glabrata* by covalent chromatography and characterization of its active centre by using 2,2′-dipyridyl disulphide as a reactivity probe 221–234
MANNING, R. *see* WALKER, J. H. 355–362
MARINKOVIĆ, D. V., PETROVIĆ, S. L., MARTINOVIĆ, J. V. & MARINKOVIĆ, J. N. Uptake of iodinated human kidney α-D-mannosidase by rat liver. Association with membrane elements and stability *in vivo* and *in vitro* 729–736
MARINKOVIĆ, J. N. *see* MARINKOVIĆ, D. V. 729–736
MARTIN, A. *see* PORTA, R. 579–583
MARTINOVIĆ, J. V. *see* MARINKOVIĆ, D. V. 729–736
MATHESON, N. R. & TRAVIS, J. Inactivation of human thrombin in the presence of human α_1-proteinase inhibitor 495–502
MATHIS, R. K. *see* BERNINGER, R. W. 95–104
MATSUBARA, H. *see* HASE, T. 55–63
MAY, R. J. & MCCARTHY, D. M. 'Particulate' hexokinase activity in rat intestine. The comparative contributions of mitochondria and brush-border membranes 189–192

MAYER, R. J. *see* PASKIN, N. 181–184; WALKER, J. H. 355–362
MCCARTHY, D. M. *see* MAY, R. J. 189–192
MEYER, T. E. *see* VAN BEEUMEN, J. 757–769
MOHOS, S. *see* SKOZA, L. 457–462
MOLE, L. E. *see* GRAND, R. J. A. 633–641
MORTON, D. J. *see* CLARKE, F. M. 797–798
MOURAY, H. *see* GAUTHIER, F. 661–665
MÜLLERTZ, S. & CLEMMENSEN, I. The primary inhibitor of plasmin in human plasma 545–553
MULLINGER, R. N. *see* HASE, T. 55–63
MURPHY, M. J., JR. ^{125}I-Labelling of erythropoietin without loss of biological activity 287–292
MURRAY, K., HUGHES, S. G., BROWN, J. S. & BRUCE, S. A. Isolation and characterization of two sequence-specific endonucleases from *Anabaena variabilis* 317–322

NOGUCHI, T., OKUNO, E. & KIDO, R. Identity of isoenzyme 1 of histidine–pyruvate aminotransferase with serine–pyruvate aminotransferase 607–613
NORRIS, R. & BROCKLEHURST, K. A convenient method of preparation of high-activity urease from *Canavalia ensiformis* by covalent chromatography and an investigation of its thiol groups with 2,2′-dipyridyl disulphide as a thiol titrant and reactivity probe 245–257
NOTARIO, V., VILLA, T. G. & VILLANUEVA, J. R. Purification of an EXO-β-D-GLUCANASE from cell-free extracts of *Candida utilis* 555–562

O'CUINN, G. *see* PIGGOTT, C. O. 715–717
OFFORD, R. E., STOREY, H. T., REES, A. R., HAYWARD, C. F., JOHNSON, W. H., PHEASEY, M. H. & WIGHTMAN, D. A. Diazoalkanes in peptide semisynthesis 480–486
OFFORD, R. E. *see also* REES, A. R. 467–479, 487–493
OGAWA, Y. *see* HATANAKA, H. 445–448
OHMIYA, N. *see* HASE, T. 55–63
OKUNO, E. *see* NOGUCHI, T. 607–613
OLD, J. M. & JONES, D. S. The aminoacylation of transfer ribonucleic acid. Inhibitory effects of some amino acid analogues with altered side chains 503–511
OLSON, J. M., SHAW, E. K. & ENGLBERGER, F. M. Comparison of bacteriochlorophyll *a*-proteins from two green bacteria 769–774
OLSON, J. M. *see also* VAN BEEUMEN, J. 757–769
OTTOLENGHI, P. *see* JENSEN, J. 815–817
OURISSON, G. *see* BOUVIER, P. 267–271

PARKHOUSE, R. M. E. *see* ABNEY, E. R. 293–299
PASKIN, N. & MAYER, R. J. Molecular weight and subunit size of fatty acid synthetase from rabbit mammary gland 181–184
PEARCE, C. J., BARNETT, J. E. G., ANTHONY, C., AKHTAR, M. & GERO, S. D. The role of the pseudo-disaccharide neamine as an intermediate in the biosynthesis of neomycin 601–606
PERHAM, R. N. *see* DANSON, M. J. 677–681
PETERS, T., JR. *see* FELDHOFF, R. C. 529–533
PETROVIĆ, S. L. *see* MARINKOVIĆ, D. V. 729–736
PETTIT, L. D. & SIDDIQUI, K. F. The proton and metal complexes of adenyl-5′-yl imidodiphosphate 169–171
PHEASEY, M. H. *see* OFFORD, R. E. 480–486
PHELPS, C. F. *see* STEVENS, R. A. J. 65–70

PIGGOTT, C. O., O'CUINN, G. & FOTTRELL, P. F. Similarities between a dipeptide hydrolase from brush-border and cytosol fractions of guinea-pig intestinal mucosa 715–717
PORTA, R., ESPOSITO, C., MARTIN, A. & DELLA PIETRA, G. Arginase from human full-term placenta 579–583
PRICE, N. C. Alternative products in the reaction of 2-nitro-5-thiocyanatobenzoic acid with thiol groups 177–180

RADDA, G. K. *see* FERGUSON, S. J. 347–353
RAO, K. K. *see* HASE, T. 55–63
REES, A. R. & OFFORD, R. E. The preparation of protected fragments of lysozyme for semisynthesis 467–479
REES, A. R. & OFFORD, R. E. The semisynthesis of portions of hen's-egg lysozyme by fragment condensation 487–493
REES, A. R. *see also* OFFORD, R. E. 480–486
REID, K. B. M. *see* BRODSKY-DOYLE, B. 279–286
REIMANN, E. M. *see* SUGDEN, P. H. 409–422
ROHMER, M. *see* BOUVIER, P. 267–271
ROSSI, A. *see* GIGLIO, J. R. 29–33

SAY, J. C. *see* GIGLIO, J. R. 29–33
SHAPOT, V. S. *see* LICHTENSTEIN, A. V. 783–794
SHAW, E. K. *see* OLSON, J. M. 769–774; VAN BEEUMEN, J. 757–769
SHIPTON, M., STUCHBURY, T. & BROCKLEHURST, K. 4-Chloro-7-nitrobenzo-2-oxa-1,3-diazole as a reactivity probe for the investigation of the thiol proteinases. Evidence that ficin and bromelain may lack carboxyl groups conformationally equivalent to that of aspartic acid-158 of papain 235–244
SIDDIQUI, K. F. *see* PETTIT, L. D. 169–171
SKOZA, L. & MOHOS, S. Stable thiobarbituric acid chromophore with dimethyl sulphoxide. Application to sialic acid assay in analytical de-*O*-acetylation 457–462
SPRINGALL, J. *see* BRITTAIN, T. 811–813
SRIVASTAVA, P. N. *see* YANG, C.-H. 133–142
STALMANS, W. *see* LEDERER, B. 689–695
STEVENS, R. A. J. & PHELPS, C. F. Studies on ligand binding to bovine liver uridine diphosphate glucose pyrophosphorylase 65–70
STORER, A. C. & CORNISH-BOWDEN, A. Concentration of $MgATP^{2-}$ and other ions in solution. Calculation of the true concentrations of species present in mixtures of associating ions 1–5
STORER, A. C. & CORNISH-BOWDEN, A. Kinetics of rat liver glucokinase. Co-operative interactions with glucose at physiologically significant concentrations 7–14
STORER, A. C. *see also* BOTTOMLEY, R. C. 667–676
STOREY, H. T. *see* OFFORD, R. E. 480–486
STUCHBURY, T. *see* SHIPTON, M. 235–244
SUBBA RAO, P. V. *see* VEERASWAMY, M. 807–809
SUGDEN, P. H. & CORBIN, J. D. Adenosine 3′:5′-cyclic monophosphate-binding proteins in bovine and rat tissues 423–437
SUGDEN, P. H., HOLLADAY, L. A., REIMANN, E. M. & CORBIN, J. D. Purification and characterization of the catalytic subunit of adenosine 3′:5′-cyclic monophosphate-dependent protein kinase from bovine liver 409–422
SUTHERLAND, I. W. *see* DAVIDSON, I. W. 707–713

TARR, G. E. & FITCH, W. M. Amino acid sequence of cytochrome *c* from *Tetrahymena pyriformis* Phenoset A 193–199
TAYLOR, R. S. & WEEDS, A. G. The magnesium-ion-dependent adenosine triphosphatase of bovine cardiac myosin and its subfragment-1 301–315
TENENHOUSE, H. S. & GOLD, R. J. M. Loss of a homologous group of proteins in a dominantly inherited ectodermal malformation 149–157
TENENHOUSE, H. S. *see also* GOLD, R. J. M. 157–160
THOMA, J. A. *see* ALLEN, J. D. 105–120, 121–131
THOMSON, A. J. *see* BRITTAIN, T. 811–813
TRAVIS, J. *see* MATHESON, N. R. 495–502
TRAYER, I. P. *see* BOTTOMLEY, R. C. 667–676
TRENTHAM, D. R. *see* ARMSTRONG, J. MCD. 513–527
TURNER, A. J. & HICK, P. E. Metabolism of daunorubicin by a barbiturate-sensitive aldehyde reductase from rat liver 819–822

VAN BAELEN, H. *see* BOUILLON, R. 463–466
VAN BEEUMEN, J., AMBLER, R. P., MEYER, T. E., KAMEN, M. D., OLSON, J. M. & SHAW, E. K. The amino acid sequences of the cytochromes *c*-555 from two green sulphur bacteria of the genus *Chlorobium* 757–769
VARIMO, K. & LONDESBOROUGH, J. Solubilization and other studies on adenylate cyclase of baker's yeast 363–370
VEERASWAMY, M., ABITHA DEVI, N., KRISHNAN KUTTY, R. & SUBBA RAO, P. V. Conversion of (±)-synephrine into *p*-hydroxyphenylacetaldehyde by *Arthrobacter synephrinum*. A novel enzymic reaction 807–809
VERNON, R. G. *see* CHRISTIE, W. W. 571–577
VILLA, T. G. *see* NOTARIO, V. 555–562
VILLANUEVA, J. R. *see* NOTARIO, V. 555–562

WALKER, J. H., BETTS, S. A., MANNING, R. & MAYER, R. J. Absorption of antisera for studies on specific enzyme turnover 355–362
WEEDS, A. G. *see* TAYLOR, R. S. 301–315
WHITEHEAD, E. P. Simplifications of the derivations and forms of steady-state equations for non-equilibrium random substrate-modifier and allosteric enzyme mechanisms 449–456
WIGHTMAN, D. A. *see* OFFORD, R. E. 480–486
WILKINSON, J. M. *see* GRAND, R. J. A. 633–641
WINZOR, D. J. *see* DREWE, R. H. 737–741
WOOD, E. J. *see* BANNISTER, J. V. 35–42
WOOSTER, M. S. & WRIGGLESWORTH, J. M. Modification of glyceraldehyde 3-phosphate dehydrogenase activity by adsorption on phospholipid vesicles 627–631
WOOTTON, J. C., KAVANAGH, J. P., BARON, A. J. & LOVETT, M. G. Re-invesigation of the effects of L-glutamine and L-asparagine on the *Neurospora crassa* NADP-specific glutamate dehydrogenase 803–806
WRIGGLESWORTH, J. M. *see* WOOSTER, M. S. 627–631

YANG, C.-H. & SRIVASTAVA, P. N. Purification and properties of arylsulphatase A from rabbit testis 133–142
YATES, D. W. & DUANCE, V. C. The binding of nucleotides and bivalent cations to the calcium-and-magnesium ion-dependent adenosine triphosphatase from rabbit muscle sarcoplasmic reticulum 719–728
YOSHIDA, A. *see* BEUTLER, E. 541–543; LEE, J. E. S. 535–539

Index of Subjects

Acetyl-coenzyme A carboxylase, absorption procedure for the preparation of monospecific antisera for, and other specific enzymes for studies on enzyme turnover (Walker, J. H., Betts, S. A., Manning, R. & Mayer, R. J.) 355–362

O-Acetylneuraminate, formation of a stable chromophore of thiobarbituric acid in the presence of dimethyl sulphoxide and its application to the assay of, in analytical de-*O*-acetylation of glycoproteins (Skoza, L. & Mohos, S.) 457–462

Actin, chick-embryo-fibroblast, structure of chicken muscle actin and (Anderson, P. J.) 185–187

Actin, identification of, and other proteins and their subunits of pig kidney-cortex microvillus membrane after sodium dodecyl sulphate/polyacrylamide-gel electrophoresis (Booth, A. G. & Kenny, A. J.) 395–407

Actin, skeletal-muscle, ox, formation of paracrystals during the binding of fructose diphosphate aldolase to filaments containing (Clarke, F. M. & Morton, D. J.) 797–798

Adenosine 3′:5′-cyclic monophosphate-binding proteins, properties of, of various ox and rat tissues (Sugden, P. H. & Corbin, J. D.) 423–437

Adenosine diphosphate, role of lipid in the binding of, to dogfish rectal-gland sodium-plus-potassium ion-dependent adenosine triphosphatase (Jensen, J. & Ottolenghi, P.) 815–817

Adenosine triphosphatase, calcium-plus-magnesium ion-dependent, sarcoplasmic-reticulum, skeletal-muscle, rabbit, binding of nucleotides and bivalent cations to (Yates, D. W. & Duance, V. C.) 719–728

Adenosine triphosphatase, magnesium ion-dependent, properties of the activity of, of ox heart-muscle myosin and of its subfragment-1 (Taylor, R. S. & Weeds, A. G.) 301–315

Adenosine triphosphatase, myosin, skeletal-muscle, rabbit, new method for quantitative affinity chromatography and its application to the study of (Bottomley, R. C., Storer, A. C. & Trayer, I. P.) 667–676

Adenosine triphosphatase, sodium-plus-potassium ion-dependent, rectal-gland, dogfish, role of lipid in the binding of adenosine diphosphate to (Jensen, J. & Ottolenghi, P.) 815–817

Adenosine triphosphatase, submitochondrial-particle, heart, ox, reaction of 4-chloro-7-nitrobenzofurazan with (Ferguson, S. J., Lloyd, W. J. & Radda, G. K.) 347–353

Adenosine triphosphate, calculation of the true concentrations of different species present in mixtures of magnesium ions and (Storer, A. C. & Cornish-Bowden, A.) 1–5

Adenylate cyclase, baker's-yeast, solubilization and other studies on (Varimo, K. & Londesborough, J.) 363–370

Adenyl-5′-yl imidodiphosphate, formation constants of complexes of, with protons and bivalent metal ions (Pettit, L. D. & Siddiqui, K. F.) 169–171

Adipose tissue, parametrial, rat, biosynthesis of triacylglycerols in homogenates of (Christie, W. W., Hunter, M. L. & Vernon, R. G.) 571–577

Albumins, serum, determination of the number and relative positions of tryptophan residues in, from various animal species (Feldhoff, R. C. & Peters, T., Jr.) 529–533

Aldehyde reductase, barbiturate-sensitive, liver, rat, reduction of daunorubicin catalysed by (Turner, A. J. & Hick, P. E.) 819–822

Aldolase, *see* Fructose diphosphate aldolase

Alginate lyase, purification and properties of, from a marine *Pseudomonas* sp. (Davidson, I. W., Sutherland, I. W. & Lawson, C. J.) 707–713

Alkaline phosphatase, *see* Phosphatase, alkaline

Alkaline phosphodiesterase, *see* Phosphodiesterase, alkaline

Amino acid compositions, method for calculating the relative contents of proteins in the major components of keratin from (Gold, R. J. M., Tenenhouse, H. S. & Adler, L. S.) 157–160

Amino acid residues, conformationally adjacent, delineation of the third antigenic site of lysozyme by application of a novel 'surface-simulation' synthetic approach directly linking, forming the site (Lee, C.-L. & Atassi, M. Z.) 89–93

Amino acid sequence of *Bacillus stearothermophilus* four-iron–four-sulphur ferredoxin (Hase, T., Ohmiya, N., Matsubara, H., Mullinger, R. N., Rao, K. K. & Hall, D. O.) 55–63

Amino acid sequence of rabbit heart-muscle troponin I (Grand, R. J. A., Wilkinson, J. M. & Mole, L. E.) 633–641

Amino acid sequence of *Tetrahymena pyriformis* (Phenoset A) cytochrome *c* (Tarr, G. E. & Fitch, W. M.) 193–199

Amino acid sequences, comparative data for, of adult major and minor β chains of haemoglobin from the two mouse species *Mus musculus* and *Mus cervicolor* (Gilman, J. G.) 43–53

Amino acid sequences of the cytochromes *c*-555 from two *Chlorobium* spp. (Van Beeumen, J., Ambler, R. P., Meyer, T. E., Kamen, M. D., Olson, J. M. & Shaw, E. K.) 757–769

Amino acids, inhibitory effects *in vitro* of some analogues of, on the aminoacylation of transfer ribonucleic acids from *Escherichia coli* and rat liver (Old, J. M. & Jones, D. S.) 503–511

Amino groups, unmasking of, of histones in chromatin at high pH (Malchy, B. L. & Kaplan, H.) 173–175

Aminoacyl-transfer ribonucleic acids, *Escherichia coli* and rat liver, inhibitory effects *in vitro* of some amino acid analogues on the formation of (Old, J. M. & Jones, D. S.) 503–511

α-Amylases, *Bacillus amyloliquefaciens* and *Aspergillus oryzae*, application of a computer model for the generation of optimum subsite maps for (Allen, J. D. & Thoma, J. A.) 121–131

Anabaena variabilis, isolation and characterization of two nucleotide-sequence-specific endodeoxyribonucleases from (Murray, K., Hughes, S. G., Brown, J. S. & Bruce, S. A.) 317–322

Antibody, anti-dinitrophenyl, rabbit, accessibility of protein-bound dinitrophenyl groups to univalent fragments of (Knight, C. G. & Green, N. M.) 323–333

Anti-dinitrophenyl antibody, rabbit, accessibility of protein-bound dinitrophenyl groups to univalent fragments of (Knight, C. G. & Green, N. M.) 323–333
Antisera, monospecific, absorption procedure for the preparation of, for studies on the turnover of specific enzymes (Walker, J. H., Betts, S. A., Manning, R. & Mayer, R. J.) 355–362
α-1-Antitrypsin, isolation and characterization of, from rhesus-monkey serum (Berninger, R. W. & Mathis, R. K.) 95–104
Apomyoglobin, sperm-whale, kinetics and mechanism of the recombination reaction between, and haemin (Adams, P. A.) 371–376
Arginase, purification and properties of, from full-term human placenta (Porta, R., Esposito, C., Martin, A. & Della Pietra, G.) 579–583
Arthrobacter synephrinum, conversion of (±)-synephrine into *p*-hydroxyphenylacetaldehyde catalysed by an enzyme from (Veeraswamy, M., Abitha Devi, N., Krishnan Kutty, R. & Subba Rao, P. V.) 807–809
Artichoke (*Helianthus tuberosus*) tubers, stability, polyadenylic acid content and ribonucleoprotein form of nuclear ribonucleic acid from (Chapman, K. S. R. & Ingle, J.) 585–600
Arylsulphatase A, purification and properties of, from rabbit testis (Yang, C.-H. & Srivastava, P. N.) 133–142
Arylsulphatase, *Charonia lampas*, substrate specificity towards carbohydrate sulphate esters of, and glycosulphatase (Hatanaka, H., Ogawa, Y. & Egami, F.) 445–448
L-Asparagine, effects of L-glutamine and, on the activity of *Neurospora crassa* nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase (Wootton, J. C., Kavanagh, J. P., Baron, A. J. & Lovett, M. G., 803–806
Aspartic acid residue 158, papain, use of 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole as a reactivity probe to provide evidence that ficin and bromelain may lack carboxyl groups conformationally equivalent to (Shipton, M., Stuchbury, T. & Brocklehurst, K.) 235–244
Aspergillus oryzae, application of a computer model for the generation of optimum subsite maps for α-amylases from *Bacillus amyloliquefaciens* and (Allen, J. D. & Thoma, J. A.) 121–131

Bacillus amyloliquefaciens, application of a computer model for the generation of optimum subsite maps for α-amylases from *Aspergillus oryzae* and (Allen, J. D. & Thoma, J. A.) 121–131
Bacillus stearothermophilus, amino acid sequence of a four-iron–four-sulphur ferredoxin from (Hase, T., Ohmiya, N., Matsubara, H., Mullinger, R. N., Rao, K. K. & Hall, D. O.) 55–63
Bacteriochlorophyll *a*, comparison of the complexes of protein with, of two *Chlorobium* spp. (Olson, J. M., Shaw, E. K. & Englberger, F. M.) 769–774
Bacteriophage PM2, effects of secondary structure on the tertiary structure of deoxyribonucleic acids of bacteriophage φX174 and (Campbell, A. M.) 615–620
Bacteriophage φX174, effects of secondary structure on the tertiary structure of deoxyribonucleic acids of bacteriophage PM2 and (Campbell, A. M.) 615–620
Bean, jack (*Canavalia ensiformis*), preparation by covalent chromatography of high-activity urease from, and use of 2,2′-dipyridyl disulphide as a thiol titrant and reactivity probe in an investigation of its thiol groups (Norris, R. & Brocklehurst, K.) 245–257
Bile pigments, model for the stereospecific degradation of haem to, *in vivo* and *in vitro* (Brown, S. B.) 23–27
Bilirubin, model for the stereospecific degradation of haem to biliverdin and, *in vivo* and *in vitro* (Brown, S. B.) 23–27
Biliverdin, model for the stereospecific degradation of haem to, and bilirubin *in vivo* and *in vitro* (Brown, S. B.) 23–27
Bone matrix, ox, comparison of the compositions of, and tendon with particular reference to the content of glycoproteins (Herring, G. M.) 749–755
Brain, rat, actions of snake venom, phospholipase A and trypsin *in vitro* on myelin purified from (Banik, N. L., Gohil, K. & Davison, A. N.) 273–277
Bromelain, use of 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole as a reactivity probe to provide evidence that ficin and, may lack carboxyl groups conformationally equivalent to aspartic acid residue 158 of papain (Shipton, M., Stuchbury, T. & Brocklehurst, K.) 235–244
Brush border, comparative contributions of mitochondria and of membranes of, to the activity of 'particulate' hexokinase in rat small intestine (May, R. J. & McCarthy, D. M.) 189–192
Brush-border fraction, mucosal, small-intestinal, guinea-pig, purification of a dipeptide hydrolase from, and similarity of its properties to those of an enzyme isolated from the cytosol fraction (Piggott, C. O., O'Cuinn, G. & Fottrell, P. F.) 715–717
Brush border, kidney-cortex, identification of the proteins and their subunits of the microvillus membrane of, after sodium dodecyl sulphate/polyacrylamide-gel electrophoresis (Booth, A. G. & Kenny, A. J.) 395–407

Calcium ions, binding of nucleotides and, to rabbit skeletal-muscle sarcoplasmic-reticulum calcium-plus-magnesium ion-dependent adenosine triphosphatase (Yates, D. W. & Duance, V. C.) 719–728
Canavalia ensiformis, *see* Bean, jack
Candida utilis C.E.C.T. 1061, purification and properties of exo-β-D-glucanase from cell-free extracts of (Notario, V., Villa, T. G. & Villanueva, J. R.) 555–562
Carbohydrate groups, location of, of chicken-egg ovomucoid (Beeley, J. G.) 335–345
Carbohydrate sulphate esters, substrate specificity of *Charonia lampas* arylsulphatase and glycosulphatase towards (Hatanaka, H., Ogawa, Y. & Egami, F.) 445–448
Carboxymethylation, effects of, of methionine residues in ox lutropin on the binding activity with receptor sites and interactions between subunits (Cheng, K.-W.) 71–77
Carboxymethylation, location of methionine residues in ox lutropin that are preferentially modified by (Cheng, K.-W.) 79–87
Cations, bivalent, binding of nucleotides and, to rabbit skeletal-muscle sarcoplasmic-reticulum calcium-plus-magnesium ion-dependent adenosine triphosphatase (Yates, D. W. & Duance, V. C.) 719–728

Cebus monkeys (*Cebus apella* and *Cebus albifrons*), comparison of the immunological and electrophoretic properties of plasma 25-hydroxycholecalciferol-binding proteins from, and from various other monkeys and primates (Bouillon, R., Van Baelen, H. & De Moor, P.) 463–466

Charonia lampas, substrate specificity towards carbohydrate sulphate esters of arylsulphatase and glycosulphatase from (Hatanaka, H., Ogawa, Y. & Egami, F.) 445–448

Chick-embryo fibroblasts, structure of actin from chicken muscle and from (Anderson, P. J.) 185–187

Chicken egg, location of the carbohydrate groups of ovomucoid from (Beeley, J. G.) 335–345

Chicken egg, preparation of protected fragments of lysozyme from, for the semisynthesis of modified polypeptides (Rees, A. R. & Offord, R. E.) 467–479

Chicken egg, semisynthesis of portions of lysozyme from, by condensation of protected fragments (Rees, A. R. & Offord, R. E.) 487–493

Chicken, structure of actin from chick-embryo fibroblasts and from muscle from (Anderson, P. J.) 185–187

Chlorobium spp., two, amino acid sequences of the cytochromes *c*-555 from (Van Beeumen, J., Ambler, R. P., Meyer, T. E., Kamen, M. D., Olson, J. M. & Shaw, E. K.) 757–769

Chlorobium spp., two, comparison of the bacteriochlorophyll *a*–protein complexes of (Olson J. M., Shaw, E. K. & Englberger, F. M.) 769–774

4-Chloro-7-nitrobenzofurazan, reaction of, with the adenosine triphosphatase of ox heart submitochondrial particles (Ferguson, S. J., Lloyd, W. J. & Radda, G. K.) 347–353

4-Chloro-7-nitrobenzofurazan, use of, as a reactivity probe to provide evidence that ficin and bromelain may lack carboxyl groups conformationally equivalent to aspartic acid residue 158 of papain (Shipton, M., Stuchbury, T. & Brocklehurst, K.) 235–244

4-Chloro-7-nitrobenzo-2-oxa-1,3-diazole, *see* 4-Chloro-7-nitrobenzofurazan

5-Chloro-4-oxopentanoate, identification of the active-site cysteine residue of rabbit skeletal-muscle pyruvate kinase as the reactive group alkylated by (Chalkley, R. A. & Bloxham, D. P.) 213–219

5-Chloro-4-oxopentanoate, selective modification of rabbit skeletal-muscle pyruvate kinase by (Bloxham, D. P. & Chalkley, R. A.) 201–211

Chlorophyll, bacterio-, *see* Bacteriochlorophyll

Cholecalciferol, comparison of the immunological and electrophoretic properties of plasma binding proteins for, and related substances from cebus monkeys and from various other monkeys and primates (Bouillon, R., Van Baelen, H. & De Moor, P.) 463–466

Chromatin, unmasking of amino groups of histones in, at high pH (Malchy, B. L. & Kaplan, H.) 173–175

Chromosome, effects of secondary structure on the tertiary structure of deoxyribonucleic acids and their relevance during the construction of models of the structure of (Campbell, A. M.) 615–620

Cibacron Blue F3GA, interaction of, with pigeon liver nicotinamide–adenine dinucleotide kinase (Apps, D. K. & Gleed, C. D.) 441–443

Complement, human, circular-dichroism and electron-microscopy studies of subcomponent C1q of component C1 of, before and after limited proteolysis by pepsin (Brodsky-Doyle, B., Leonard, K. R. & Reid, K. B. M.) 279–286

Component C1, complement, human, circular-dichroism and electron-microscopy studies of subcomponent C1q of, before and after limited proteolysis by pepsin (Brodsky-Doyle, B., Leonard, K. R. & Reid, K. B. M.) 279–286

Computer model, application of, for the generation of optimum subsite maps for *Bacillus amyloliquefaciens* and *Aspergillus oryzae* α-amylases (Allen, J. D. & Thoma, J. A.) 121–131

Computer model, development of, for the generation of optimum subsite maps for depolymerases (Allen, J. D. & Thoma, J. A.) 105–120

Creatine kinase, skeletal-muscle, rabbit, alternative products in the reaction of 2-nitro-5-thiocyanatobenzoic acid with thiol groups of (Price, N. C.) 177–180

Cyclic adenosine 3′:5′-monophosphate, *see* Adenosine 3′:5′-cyclic monophosphate

Cysteine residue, active-site, identification of, of rabbit skeletal-muscle pyruvate kinase as the reactive group alkylated by 5-chloro-4-oxopentanoate (Chalkley, R. A. & Bloxham, D. P.) 213–219

Cysteine transfer ribonucleic acid, baker's-yeast, application of a spectrophotometric method to the determination of the composition of oligonucleotides obtained by ribonuclease digestion of (Holness, N. J. & Atfield, G.) 15–22

Cytochrome *c*, [methionine sulphoxide]-, kinetic studies on (Brittain, T. & Greenwood, C.) 621–626

Cytochrome *c* oxidase, absorption procedure for the preparation of monospecific antisera for, and other specific enzymes for studies on enzyme turnover (Walker, J. H., Betts, S. A., Manning, R. & Mayer, R. J.) 355–362

Cytochrome *c* oxidase, heart-muscle, ox, evidence for haem–haem interactions obtained during low-temperature studies of, by using magnetic circular dichroism (Brittain, T., Springall, J., Greenwood, C. & Thomson, A. J.) 811–813

Cytochrome *c*, *Tetrahymena pyriformis* (Phenoset A), amino acid sequence of (Tarr, G. E. & Fitch, W. M.) 193–199

Cytochromes *c*-555, *Chlorobium* spp., two, amino acid sequences of (Van Beeumen, J., Ambler, R. P., Meyer, T. E., Kamen, M. D., Olson, J. M. & Shaw, E. K.) 757–769

Cytosol, mucosal, small-intestinal, guinea-pig, similarity of the properties of a dipeptide hydrolase purified from the brush-border fraction to those of an enzyme isolated from (Piggott, C. O., O'Cuinn, G. & Fottrell, P. F.) 715–717

Cytosol, role of, in governing the regulation of the transcription of the gene deoxyribonucleic acids in the biosynthesis of messenger ribonucleic acids (Lichtenstein, A. V. & Shapot, V. S.) 783–794

Daunorubicin, reduction of, catalysed by a rat liver barbiturate-sensitive aldehyde reductase (Turner, A. J. & Hick, P. E.) 819–822

Deoxyribonucleases, endo-, nucleotide-sequence-specific, two, isolation and characterization of, from *Anabaena variabilis* (Murray, K., Hughes, S. G., Brown, J. S. & Bruce, S. A.) 317–322

Deoxyribonucleic acids, bacteriophage, effects of secondary structure on the tertiary structure of (Campbell, A. M.) 615–620

Deoxyribonucleic acids, gene, model for the cytosol-governed regulation of the transcription of, in the biosynthesis of messenger ribonucleic acids (Lichtenstein, A. V. & Shapot, V. S.) 783–794

Depolymerases, development of a computer model for the generation of optimum subsite maps for (Allen, J. D. & Thoma, J. A.) 105–120

Diazoalkanes, use of, as protecting agents in the semisynthesis of polypeptides (Offord, R. E., Storey, H. T., Rees, A. R., Hayward, C. F., Johnson, W. H., Pheasey, M. H. & Wightman, D. A.) 480–486

4,4-Dimethyl sterols, identification of 4α-methyl sterols and, in *Methylococcus capsulatus* as $\Delta^{8(14)}$-steroids (Bouvier, P., Rohmer, M., Benveniste, P. & Ourisson, G.) 267–271

Dinitrophenyl groups, protein bound, accessibility of, to univalent fragments of rabbit anti-dinitrophenyl antibody (Knight, C. G. & Green, N. M.) 323–333

N-(*N*-Dinitrophenylaminoalkyl)maleimides, reaction of, with the thiol group of ox mercaptalbumin (Knight, C. G. & Green, N. M.) 323–333

Dipeptide hydrolase, purification of, from guinea-pig small-intestinal mucosal brush-border fraction and similarity of its properties to those of an enzyme isolated from the cytosol fraction (Piggott, C. O., O'Cuinn, G. & Fottrell, P. F.) 715–717

2,2′-Dipyridyl disulphide, use of, as a reactivity probe in the characterization of the active centre of fully active ficin prepared from *Ficus glabrata* (Malthouse, J. P. G. & Brocklehurst, K.) 221–234

2,2′-Dipyridyl disulphide, use of, as a thiol titrant and reactivity probe in an investigation of the thiol groups of high-activity urease prepared from jack-bean meal (Norris, R. & Brocklehurst, K.) 245–257

Dogfish, role of lipid in the binding of adenosine diphosphate to sodium-plus-potassium ion-dependent adenosine triphosphatase from the rectal gland of (Jensen, J. & Ottolenghi, P.) 815–817

Egg, chicken, location of the carbohydrate groups of ovomucoid from (Beeley, J. G.) 335–345

Egg, chicken, preparation of protected fragments of lysozyme from, for the semisynthesis of modified polypeptides (Rees, A. R. & Offord, R. E.) 467–479

Egg, chicken, semisynthesis of portions of lysozyme from, by condensation of protected fragments (Rees, A. R. & Offord, R. E.) 487–493

Endodeoxyribonucleases, nucleotide-sequence-specific, two, isolation and characterization of, from *Anabaena variabilis* (Murray, K., Hughes, S. G., Brown, J. S. & Bruce, S. A.) 317–322

Enzymes, development of a computer model for the generation of optimum subsite maps for (Allen, J. D. & Thoma, J. A.) 105–120

Enzymes, practical difficulties with mechanisms that contain repeated rate constants in algebraic methods for deriving steady-state rate equations for reactions catalysed by (Cornish-Bowden, A.) 167

Enzymes, pre-eminence of $k_{cat.}$ in the manifestation of optimum activity delineated by using the Briggs-Haldane two-step irreversible kinetic model for reactions catalysed by (Brocklehurst, K. & Cornish-Bowden, A.) 165–166

Enzymes, simplifications of the derivations and forms of steady-state equations for non-equilibrium random substrate-modifier and allosteric mechanisms for reactions catalysed by (Whitehead, E. P.) 449–456

Enzymes, specific, absorption procedure for the preparation of monospecific antisera for studies on the turnover of (Walker, J. H., Betts, S. A., Manning, R. & Mayer, R. J.) 355–362

Erythrocruorin, isolation, characterization and oxygen equilibrium of, from the polychaete worm *Eunice aphroditois* (Bannister, J. V., Bannister, W. H., Anastasi, A. & Wood, E. J.) 35–42

Erythropoietin, sheep and human, ^{125}I-labelling of, without loss of biological activity (Murphy, M. J., Jr.) 287–292

Escherichia coli, inhibitory effects *in vitro* of some amino acid analogues on the aminoacylation of transfer ribonucleic acids from rat liver and (Old, J. M. & Jones, D. S.) 503–511

Escherichia coli K12, evidence for the presence of two lipoic acid residues per chain of lipoate acetyltransferase in the pyruvate dehydrogenase multienzyme complex of (Danson, M. J. & Perham, R. N.) 677–681

Eunice aphroditois, *see* Polychaete worm

Exo-β-D-glucanase, purification and properties of, from *Candida utilis* C.E.C.T. 1061 (Notario, V., Villa, T. G. & Villanueva, J. R.) 555–562

Fatty acid synthetase, mammary-gland, rabbit, molecular weight and subunit size of (Paskin, N. & Mayer, R. J.) 181–184

Ferredoxin, four-iron–four-sulphur, *Bacillus stearothermophilus*, amino acid sequence of (Hase, T., Ohmiya, N., Matsubara, H., Mullinger, R. N., Rao, K. K. & Hall, D. O.) 55–63

Fibrinolysis, properties of the primary inhibitor of plasmin in human plasma and their relevance to (Müllertz, S. & Clemmensen, I.) 545–553

Fibroblasts, chick-embryo, structure of actin from chicken muscle and from (Anderson, P. J.) 185–187

Ficin, preparation by covalent chromatography of, from *Ficus glabrata* and characterization of its active centre by the use of 2,2′-dipyridyl disulphide as a reactivity probe (Malthouse, J. P. G. & Brocklehurst, K.) 221–234

Ficin, use of 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole as a reactivity probe to provide evidence that bromelain and, may lack carboxyl groups conformationally equivalent to aspartic acid residue 158 of papain (Shipton, M., Stuchbury, T. & Brocklehurst, K.) 235–244

Ficus glabrata, preparation by covalent chromatography of fully active ficin from, and characterization of its active centre by the use of 2,2′-dipyridyl disulphide as a reactivity probe (Malthouse, J. P. G. & Brocklehurst, K.) 221–234

Follicle-stimulating hormone, *see* Follitropin
Follitropin, purification and properties of, from ox anterior pituitary gland (Cheng, K.-W.) 651–659
Fructose diphosphate aldolase, skeletal-muscle, ox, formation of paracrystals during the binding of, to filaments containing actin (Clarke, F. M. & Morton, D. J.) 797–798

Gastrin, inhibitory effects *in vitro* of analogues of the *C*-terminal tetrapeptide of, on the aminoacylation of rat liver and *Escherichia coli* transfer ribonucleic acids (Old, J. M. & Jones, D. S.) 503–511
Gel filtration, degree of removal of bound ligands from macromolecules by (Dixon, H. B. F.) 161–162
Genes, model for the cytosol-governed regulation of the transcription of the deoxyribonucleic acids of, in the biosynthesis of messenger ribonucleic acids (Lichtenstein, A. V. & Shapot, V. S.) 783–794
α_1-Globulin, isolation and characterization as α-1-antitrypsin of, from rhesus-monkey serum (Berninger, R. W. & Mathis, R. K.) 95–104
β-D-Glucanase, exo-, purification and properties of, from *Candida utilis* C.E.C.T. 1061 (Notario, V., Villa, T. G. & Villanueva, J. R.) 555–562
Glucokinase, liver, rat, kinetics of the reaction catalysed by, showing co-operative interactions of the enzyme with glucose at physiologically significant concentrations (Storer, A. C. & Cornish-Bowden, A.) 7–14
Glucose, kinetics of the reaction catalysed by rat liver glucokinase showing co-operative interactions of the enzyme with, at physiologically significant concentrations (Storer, A. C. & Cornish-Bowden, A.) 7–14
Glucose 6-phosphate dehydrogenase, characterization of the activity of, in rat liver mitochondria (Grunwald, M. & Hill, H. Z.) 683–687
Glutamate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-specific, *Neurospora crassa*, effects of L-glutamine and L-asparagine on the activity of (Wootton, J. C., Kavanagh, J. P., Baron, A. J. & Lovett, M. G.) 803–806
L-Glutamine, effects of L-asparagine and, on the activity of *Neurospora crassa* nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase (Wootton, J. C., Kavanagh, J. P., Baron, A. J. & Lovett, M. G.) 803–806
Glutathione–insulin transhydrogenase, membrane-bound, microsomal, liver, rat, activity of, and comparison of its properties with those of protein disulphide-isomerase (Ibbetson, A. L. & Freedman, R. B.) 377–384
Glutathione–insulin transhydrogenase, microsomal, liver, ox, differences between the activities of, and of protein disulphide-isomerase (Hawkins, H. C. & Freedman, R. B.) 385–393
D-Glyceraldehyde 3-phosphate dehydrogenase, skeletal-muscle, pig, reactions of D-glyceraldehyde 3-phosphate with thiols and with the holoenzyme of, and of inorganic phosphate with the acyl-holoenzyme (Armstrong, J. McD. & Trentham, D. R.) 513–527
Glyceraldehyde 3-phosphate dehydrogenase, skeletal-muscle, rabbit, modification of the activity of, by adsorption on phospholipid vesicles (Wooster, M. S. & Wrigglesworth, J. M.) 627–631
D-Glyceraldehyde 3-phosphate, reactions of, with thiols and with the holoenzyme of pig skeletal-muscle D-glyceraldehyde 3-phosphate dehydrogenase (Armstrong, J. McD. & Trentham, D. R.) 513–527
Glycogen phosphorylase, kinetic properties of, from human liver and assay of its activity in biopsy specimens (Lederer, B. & Stalmans, W.) 689–695
Glycoprotein (ovomucoid), chicken-egg, location of the carbohydrate groups of (Beeley, J. G.) 335–345
Glycoproteins, comparison of the compositions of ox bone matrix and tendon with particular reference to the content of (Herring, G. M.) 749–755
Glycoproteins, formation of a stable chromophore of thiobarbituric acid in the presence of dimethyl sulphoxide and its application to the assay of sialic acid in analytical de-*O*-acetylation of (Skoza, L. & Mohos, S.) 457–462
Glycosaminoglycans, other, method for the determination of hyaluronate in the presence of, and its application to human intervertebral disc (Hardingham, T. E. & Adams, P.) 143–147
Glycosulphatase, *Charonia lampas*, substrate specificity towards carbohydrate sulphate esters of, and arylsulphatase (Hatanaka, H., Ogawa, Y. & Egami, F.) 445–448
Growth hormone, purification and characterization of rabbit placental lactogen and its relationship to (Bolander, F. F., Jr. & Fellows, R. E.) 775–782

Haem–haem interactions, evidence for, obtained during low-temperature studies of ox heart-muscle cytochrome *c* oxidase by using magnetic circular dichroism (Brittain, T., Springall, J., Greenwood, C. & Thomson, A. J.) 811–813
Haem, model for the stereospecific degradation of, to bile pigments *in vivo* and *in vitro* (Brown, S. B.) 23–27
Haemin, kinetics and mechanism of the recombination reaction between sperm-whale apomyoglobin and (Adams, P. A.) 371–376
Haemoglobin, extracellular, isolation, characterization and oxygen equilibrium of, from the polychaete worm *Eunice aphroditois* (Bannister, J. V., Bannister, W. H., Anastasi, A. & Wood, E. J.) 35–42
Haemoglobin, mouse, comparative data for the amino acid sequences of adult major and minor β chains of, from the two species *Mus musculus* and *Mus cervicolor* (Gilman, J. G.) 43–53
Heart muscle, ox, evidence for haem–haem interactions obtained during low-temperature studies of cytochrome *c* oxidase from, by using magnetic circular dichroism (Brittain, T., Springall, J., Greenwood, C. & Thomson, A. J.) 811–813
Heart muscle, ox, properties of the magnesium ion-dependent adenosine triphosphatase activity of myosin and of its subfragment-1 from (Taylor, R. S. & Weeds, A. G.) 301–315
Heart muscle, rabbit, amino acid sequence of troponin I from (Grand, R. J. A., Wilkinson, J. M. & Mole, L. E.) 633–641
Heart, ox, reaction of 4-chloro-7-nitrobenzofurazan with the adenosine triphosphatase of submitochondrial particles from (Ferguson, S. J., Lloyd, W. J. & Radda, G. K.) 347–353
Helianthus tuberosus, *see* Artichoke
Hen, *see* Chicken

Hexokinase, 'particulate', comparative contributions of mitochondria and brush-border membranes to the activity of, in rat small intestine (May, R. J. & McCarthy, D. M.) 189–192
Hexosaminidase A, subunit structure of, from human placenta (Beutler, E., Yoshida, A., Kuhl, W. & Lee, J. E. S.) 541–543
Hexosaminidases A and B, purification and chemical characterization of, from human placenta (Lee, J. E. S. & Yoshida, A.) 535–539
Histidine–pyruvate aminotransferase, liver, rat, identity of isoenzyme 1 of, with serine–pyruvate aminotransferase (Noguchi, T., Okuno, E. & Kido, R.) 607–613
Histones, unmasking of amino groups of, in chromatin at high pH (Malchy, B. L. & Kaplan, H.) 173–175
Hyaluronate, method for the determination of, in the presence of other glycosaminoglycans and its application to human intervertebral disc (Hardingham, T. E. & Adams, P.) 143–147
Hydrogen ions, formation constants of complexes of adenyl-5′-yl imidodiphosphate with bivalent metal ions and (Pettit, L. D. & Siddiqui, K. F.) 169–171
Hydrolases, development of a computer model for the generation of optimum subsite maps for (Allen, J. D. & Thoma, J. A.) 105–120
25-Hydroxycholecalciferol-binding proteins, plasma, comparison of the immunological and electrophoretic properties of, from cebus monkeys and from various other monkeys and primates (Bouillon, R., Van Baelen, H. & De Moor, P.) 463–466
4-Hydroxypentanoic acid alanine thioether, synthesis and characterization of (Chalkley, R. A. & Bloxham, D. P.) 213–219
p-Hydroxyphenylacetaldehyde, conversion of (±)-synephrine into, catalysed by an enzyme from *Arthrobacter synephrinum* (Veeraswamy, M., Abitha Devi, N., Krishnan Kutty, R. & Subba Rao, P. V.) 807–809

Inorganic phosphate, *see* Phosphate, inorganic
Intervertebral disc, human, method for the determination of hyaluronate in the presence of other glycosaminoglycans and its application to (Hardingham, T. E. & Adams, P.) 143–147
Intestine, small, guinea-pig, purification of a dipeptide hydrolase from the brush-border fraction of the mucosa of, and similarity of its properties to those of an enzyme isolated from the cytosol fraction (Piggott, C. O., O'Cuinn, G. & Fottrell, P. F.) 715–717
Intestine, small, rat, comparative contributions of mitochondria and brush-border membranes to the activity of 'particulate' hexokinase in (May, R. J. & McCarthy, D. M.) 189–192
Iodine isotope (^{125}I), labelling of sheep and human erythropoietin with, without loss of biological activity (Murphy, M. J., Jr.) 287–292
Ions, associating, calculation of the true concentrations of different species present in mixtures of (Storer, A. C. & Cornish-Bowden, A.) 1–5
Ions, electrophoretic method for the study of the reversible binding of, to proteins (Drewe, R. H. & Winzor, D. J.) 737–741

Jack bean, *see* Bean, jack
Jerusalem artichoke, *see* Artichoke

Keratin, method for calculating the relative contents of proteins in the major components of, from their amino acid compositions (Gold, R. J. M., Tenenhouse, H. S. & Adler, L. S.) 157–160
Keratin, mouse, loss of a homologous group of proteins from, of animals with the dominantly inherited Naked trait (Tenenhouse, H. S. & Gold, R. J. M.) 149–157
Kidney cortex, pig, identification of the proteins and their subunits of the microvillus membrane of, after sodium dodecyl sulphate/polyacrylamide-gel electrophoresis (Booth, A. G. & Kenny, A. J.) 395–407
Kidney, human, uptake of α-D-mannosidase purified from, by rat liver and evidence for association of the enzyme with membrane elements and its stability *in vivo* and *in vitro* (Marinković, D. V., Petrović, S. L., Martinović, J. V. & Marinković, J. N.) 729–736
Kinetic model, irreversible, two-step, Briggs–Haldane, pre-eminence of $k_{cat.}$ in the manifestation of optimum enzymic activity by using (Brocklehurst, K. & Cornish-Bowden, A.) 165–166
Kinetics, non-Michaelian, simplifications of the derivations and forms of steady-state equations for, of non-equilibrium random substrate-modifier and allosteric enzyme mechanisms (Whitehead, E. P.) 449–456

Lactogen, placental, rabbit, purification and characterization of (Bolander, F. F., Jr. & Fellows, R. E.) 775–782
Lecithin, *see* Phosphatidylcholine
Ligands, binding of, to ox liver uridine diphosphate glucose pyrophosphorylase (Stevens, R. A. J. & Phelps, C. F.) 65–70
Ligands, bound, degree of removal of, from macromolecules by gel filtration (Dixon, H. B. F.) 161–162
Lipid, role of, in the binding of adenosine diphosphate to dogfish rectal-gland sodium-plus-potassium ion-dependent adenosine triphosphatase (Jensen, J. & Ottolenghi, P.) 815–817
Lipoate acetyltransferase, evidence for the presence of two lipoic acid residues per chain of, in the *Escherichia coli* K12 pyruvate dehydrogenase multienzyme complex (Danson, M. J. & Perham, R. N.) 677–681
Lipoic acid residues, two, evidence for the presence of, per chain of lipoate acetyltransferase in the *Escherichia coli* K12 pyruvate dehydrogenase multienzyme complex (Danson, M. J. & Perham, R. N.) 677–681
Liposomes, modification of the activity of rabbit skeletal-muscle glyceraldehyde 3-phosphate dehydrogenase by adsorption on (Wooster, M. S. & Wrigglesworth, J. M.) 627–631
Liver, human, kinetic properties of glycogen phosphorylase from, and assay of its activity in biopsy specimens (Lederer, B. & Stalmans, W.) 689–695
Liver, human, purification and subunit structure of alkaline phosphatase from (Latner, A. L. & Hodson, A. W.) 697–705
Liver, ox and rat, properties of adenosine 3′:5′-cyclic monophosphate-binding proteins of, and other tissues (Sugden, P. H. & Corbin, J. D.) 423–437
Liver, ox, binding of ligands to uridine diphosphate glucose pyrophosphorylase from (Stevens, R. A. J. & Phelps, C. F.) 65–70

Liver, ox, differences between the activities of microsomal protein disulphide-isomerase and glutathione–insulin transhydrogenase in (Hawkins, H. C. & Freedman, R. B.) 385–393
Liver, ox, purification and physical properties of the catalytic subunit of adenosine 3′:5′-cyclic monophosphate-dependent protein kinase from (Sugden, P. H., Holladay, L. A., Reimann, E. M. & Corbin, J. D.) 409–422
Liver, pigeon, interaction of Cibacron Blue F3GA with nicotinamide–adenine dinucleotide kinase from (Apps, D. K. & Gleed, C. D.) 441–443
Liver, rat, activity of microsomal membrane-bound glutathione–insulin transhydrogenase in, and comparison of its properties with those of protein disulphide-isomerase (Ibbetson, A. L. & Freedman, R. B.) 377–384
Liver, rat, characterization of the activity of glucose 6-phosphate dehydrogenase in mitochondria from (Grunwald, M. & Hill, H. Z.) 683–687
Liver, rat, identity of isoenzyme 1 of histidine–pyruvate aminotransferase from, with serine–pyruvate aminotransferase (Noguchi, T., Okuno, E. & Kido, R.) 607–613
Liver, rat, inhibitory effects *in vitro* of some amino acid analogues on the aminoacylation of transfer ribonucleic acids from *Escherichia coli* and (Old, J. M. & Jones, D. S.) 503–511
Liver, rat, kinetics of the reaction catalysed by glucokinase from, showing co-operative interactions of the enzyme with glucose at physiologically significant concentrations (Storer, A. C. & Cornish-Bowden, A.) 7–14
Liver, rat, reduction of daunorubicin catalysed by a barbiturate-sensitive aldehyde reductase from (Turner, A. J. & Hick, P. E.) 819–822
Liver, rat, uptake of iodinated human kidney α-D-mannosidase by, and evidence for association of the enzyme with membrane elements and its stability *in vivo* and *in vitro* (Marinković, D. V., Petrović, S. L., Martinović, J. V. & Marinković, J. N.) 729–736
Luteinizing hormone, *see* Lutropin
Lutropin, ox, effects of carboxymethylation of methionine residues in, on the binding activity with receptor sites and interactions between subunits (Cheng, K.-W.) 71–77
Lutropin, ox, location of methionine residues in, that are preferentially modified by carboxymethylation (Cheng, K.-W.) 79–87
Lymphocytes, mouse, location of the activities of nucleotide pyrophosphatase and alkaline phosphodiesterase on the plasma membrane of (Abney, E. R., Evans, W. H. & Parkhouse, R. M. E.) 293–299
Lysozyme, chicken-egg, preparation of protected fragments of, for the semisynthesis of modified polypeptides (Rees, A. R. & Offord, R. E.) 467–479
Lysozyme, chicken-egg, semisynthesis of portions of, by condensation of protected fragments (Rees, A. R. & Offord, R. E.) 487–493
Lysozyme, delineation of the third antigenic site of, by application of a novel 'surface-simulation' synthetic approach directly linking the conformationally adjacent amino acid residues forming the site (Lee, C.-L. & Atassi, M. Z.) 89–93
Macaca mulatta, *see* Rhesus monkey
α_2-Macroglobulin, acute-phase, isolation and physicochemical properties of, from rat serum (Gauthier, F. & Mouray, H.) 661–665
α_1- and α_2-Macroglobulins, acute-phase, isolation and properties of, from rat serum (Gordon, A. H.) 643–650
Macromolecules, degree of removal of bound ligands from, by gel filtration (Dixon, H. B. F.) 161–162
Magnesium–adenosine triphosphate complex ion, calculation of the true concentrations of, and other species present in mixtures of adenosine triphosphate and magnesium ions (Storer, A. C. & Cornish-Bowden, A.) 1–5
Magnesium ions, calculation of the true concentrations of different species present in mixtures of adenosine triphosphate and (Storer, A. C. & Cornish-Bowden, A.) 1–5
Maleimides, use of, as chemical probes to measure the content of lipoic acid in the *Escherichia coli* K12 pyruvate dehydrogenase multienzyme complex (Danson, M. J. & Perham, R. N.) 677–681
Mammary gland, rabbit, molecular weight and subunit size of fatty acid synthetase from (Paskin, N. & Mayer, R. J.) 181–184
α-D-Mannosidase, kidney, human, iodinated, uptake of, by rat liver and evidence for association of the enzyme with membrane elements and its stability *in vivo* and *in vitro* (Marinković, D. V., Petrović, S. L., Martinović, J. V. & Marinković, J. N.) 729–736
Membrane, microvillus, kidney-cortex, pig, identification of the proteins and their subunits of, after sodium dodecyl sulphate/polyacrylamide-gel electrophoresis (Booth, A. G. & Kenny, A. J.) 395–407
Membrane, plasma, lymphocyte, mouse, location of the activities of nucleotide pyrophosphatase and alkaline phosphodiesterase on (Abney, E. R., Evans, W. H. & Parkhouse, R. M. E.) 293–299
Membranes, brush-border, comparative contributions of mitochondria and of, to the activity of 'particulate' hexokinase in rat small intestine (May, R. J. & McCarthy, D. M.) 189–192
Membranes, liver, rat, association of iodinated human kidney α-D-mannosidase with, after intravenous injection of the enzyme into the animals (Marinković, D. V., Petrović, S. L., Martinović, J. V. & Marinković, J. N.) 729–736
Membranes, microsomal, liver, rat, activity of glutathione–insulin transhydrogenase bound to, and comparison of its properties with those of protein disulphide-isomerase (Ibbetson, A. L. & Freedman, R. B.) 377–384
Membranes, submitochondrial-particle, heart, ox, reaction of 4-chloro-7-nitrobenzofurazan with the adenosine triphosphatase of (Ferguson, S. J., Lloyd, W. J. & Radda, G. K.) 347–353
Mercaptalbumin, ox, reaction of *N*-(*N*-dinitrophenylaminoalkyl)maleimides with the thiol group of (Knight, C. G. & Green, N. M.) 323–333
Messenger ribonucleic acid, *see* Ribonucleic acid, messenger
Metal ions, bivalent, formation constants of complexes of adenyl-5′-yl imidodiphosphate with protons and (Pettit, L. D. & Siddiqui, K. F.) 169–171

Methionine residues, effects of carboxymethylation of, in ox lutropin on the binding activity with receptor sites and interactions between subunits (Cheng, K.-W.) 71–77
[Methionine sulphoxide]cytochrome *c*, kinetic studies on (Brittain, T. & Greenwood, C.) 621–626
4α-Methyl sterols, identification of 4,4-dimethyl sterols and, in *Methylococcus capsulatus* as $\Delta^{8(14)}$-steroids (Bouvier, P., Rohmer, M., Benveniste, P. & Ourisson, G.) 267–271
Methylococcus capsulatus, identification of $\Delta^{8(14)}$-steroids in (Bouvier, P., Rohmer, M., Benveniste, P. & Ourisson, G.) 267–271
Microsomal fraction, liver, ox, differences between the activities of protein disulphide-isomerase and glutathione–insulin transhydrogenase in (Hawkins, H. C. & Freedman, R. B.) 385–393
Microsomal fraction, liver, rat, activity of membrane-bound glutathione–insulin transhydrogenase in, and comparison of its properties with those of protein disulphide-isomerase (Ibbetson, A. L. & Freedman, R. B.) 377–384
Microvillus membrane, kidney-cortex, pig, identification of the proteins and their subunits of, after sodium dodecyl sulphate/polyacrylamide-gel electrophoresis (Booth, A. G. & Kenny, A. J.) 395–407
Mitochondria, comparative contributions of brush-border membranes and of, to the activity of 'particulate' hexokinase in (May, R. J. & McCarthy, D. M.) 189–192
Mitochondria, heart, ox, reaction of 4-chloro-7-nitrobenzofurazan with the adenosine triphosphatase of submitochondrial particles from (Ferguson, S. J., Lloyd, W. J. & Radda, G. K.) 347–353
Mitochondria, liver, rat, characterization of the activity of glucose 6-phosphate dehydrogenase in (Grunwald, M. & Hill, H. Z.) 683–687
Mucosa, small-intestinal, guinea-pig, purification of a dipeptide hydrolase from the brush-border fraction of, and similarity of its properties to those of an enzyme isolated from the cytosol fraction (Piggott, C. O., O'Cuinn, G. & Fottrell, P. F.) 715–717
Muscle, chicken, structure of actin from chick-embryo fibroblasts and from (Anderson, P. J.) 185–187
Muscle, heart, ox, evidence for haem–haem interactions obtained during low-temperature studies of cytochrome *c* oxidase from, by using magnetic circular dichroism (Brittain, T., Springall, J., Greenwood, C. & Thomson, A. J.) 811–813
Muscle, heart, ox, properties of the magnesium ion-dependent adenosine triphosphatase activity of myosin and of its subfragment-1 from (Taylor, R. S. & Weeds, A. G.) 301–315
Muscle, heart, rabbit, amino acid sequence of troponin I from (Grand, R. J. A., Wilkinson, J. M. & Mole, L. E.) 633–641
Muscle, skeletal, ox, formation of paracrystals during the binding of fructose diphosphate aldolase to filaments containing actin from (Clarke, F. M. & Morton, D. J.) 797–798
Muscle, skeletal, pig, reactions of D-glyceraldehyde 3-phosphate with thiols and with the holoenzyme of D-glyceraldehyde 3-phosphate dehydrogenase from, and of inorganic phosphate with the acyl-holoenzyme (Armstrong, J. McD. & Trentham, D. R.) 513–527
Muscle, skeletal, rabbit, binding of nucleotides and bivalent cations to the calcium-plus-magnesium ion-dependent adenosine triphosphatase from the sarcoplasmic reticulum of (Yates, D. W. & Duance, V. C.) 719–728
Muscle, skeletal, rabbit, identification of the active-site cysteine residue of pyruvate kinase from, as the reactive group alkylated by 5-chloro-4-oxopentanoate (Chalkley, R. A. & Bloxham, D. P.) 213–219
Muscle, skeletal, rabbit, modification of the activity of glyceraldehyde 3-phosphate dehydrogenase from, by adsorption on phospholipid vesicles (Wooster, M. S. & Wrigglesworth, J. M.) 627–631
Muscle, skeletal, rabbit, new method for quantitative affinity chromatography and its application to the study of myosin from (Bottomley, R. C., Storer, A. C. & Trayer, I. P.) 667–676
Muscle, skeletal, rabbit, selective modification by 5-chloro-4-oxopentanoate of pyruvate kinase from (Bloxham, D. P. & Chalkley, R. A.) 201–211
Myelin, rat, purified, actions of snake venom, phospholipase A and trypsin *in vitro* on (Banik, N. L., Gohil, K. & Davison, A. N.) 273–277
Myoglobin, apo-, sperm-whale, kinetics and mechanism of the recombination reaction between, and haemin (Adams, P. A.) 371–376
Myosin, heart-muscle, ox, properties of the magnesium ion-dependent adenosine triphosphatase activity of, and of its subfragment-1 (Taylor, R. S. & Weeds, A. G.) 301–315
Myosin, skeletal-muscle, rabbit, new method for quantitative affinity chromatography and its application to the study of (Bottomley, R. C., Storer, A. C. & Trayer, I. P.) 667–676

Naja naja, *see* Snake
Neamine, role of, in the biosynthesis of neomycin in *Streptomyces* spp. (Pearce, C. J., Barnett, J. E. G., Anthony, C., Akhtar, M. & Gero, S. D.) 601–606
Neomycin, role of the pseudo-disaccharide neamine in the biosynthesis of, in *Streptomyces* spp. (Pearce, C. J., Barnett, J. E. G., Anthony, C., Akhtar, M. & Gero, S. D.) 601–606
Neurospora crassa, effects of L-glutamine and L-asparagine on the activity of nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase from (Wootton, J. C., Kavanagh, J. P., Baron, A. J. & Lovett, M. G.) 803–806
Nicotinamide–adenine dinucleotide kinase, liver, pigeon, interaction of Cibacron Blue F3GA with (Apps, D. K & Gleed, C. D.) 441–443
2-Nitro-5-thiocyanatobenzoic acid, alternative products in the reaction of, with thiol groups of proteins (Price, N. C.) 177–180
Nuclear ribonucleic acid, *see* Ribonucleic acid, nuclear
Nucleotide composition, application of a spectrophotometric method to the determination of, of oligonucleotides obtained by ribonuclease digestion of baker's-yeast cysteine transfer ribonucleic acid (Holness, N. J. & Atfield, G.) 15–22
Nucleotide pyrophosphatase, location of the activities of alkaline phosphodiesterase and, on mouse lymphocyte plasma membrane (Abney, E. R., Evans, W. H. & Parkhouse, R. M. E.) 293–299

Nucleotide sequences, specificity for, of two endodeoxyribonucleases isolated from *Anabaena variabilis* (Murray, K., Hughes, S. G., Brown, J. S. & Bruce, S. A.) 317–322
Nucleotides, binding of bivalent cations and, to rabbit skeletal-muscle sarcoplasmic-reticulum calcium-plus-magnesium ion-dependent adenosine triphosphatase (Yates, D. W. & Duance, V. C.) 719–728

Oligonucleotides, application of a spectrophotometric method to the determination of the composition of, obtained by ribonuclease digestion of baker's-yeast cysteine transfer ribonucleic acid (Holness, N. J. & Atfield, G.) 15–22
Ovalbumin, electrophoretic study of the reversible binding of inorganic phosphate to (Drewe, R. H. & Winzor, D. J.) 737–741
Ovomucoid, chicken-egg, location of the carbohydrate groups of (Beeley, J. G.) 335–345
Oxygen equilibrium, isolation, characterization and, of an extracellular haemoglobin from the polychaete worm *Eunice aphroditois* (Bannister, J. V., Bannister, W. H., Anastasi, A. & Wood, E. J.) 35–42

Papain, use of 4-chloro-7-nitrobenzo-2-oxa-1,3-diazole as a reactivity probe to provide evidence that ficin and bromelain may lack carboxyl groups conformationally equivalent to aspartic acid residue 158 of (Shipton, M., Stuchbury, T. & Brocklehurst, K.) 235–244
Peptides, use of diazoalkanes as protecting agents in the semisynthesis of (Offord, R. E., Storey, H. T., Rees, A. R., Hayward, C. F., Johnson, W. H., Pheasey, M. H. & Wightman, D. A.) 480–486
pH, high, unmasking of amino groups of histones in chromatin at (Malchy, B. L. & Kaplan, H.) 173–175
Phage, *see* Bacteriophage
Phosphatase, alkaline, purification and subunit structure of, from human liver (Latner, A. L. & Hodson, A. W.) 697–705
Phosphate, inorganic, electrophoretic study of the reversible binding of, to ovalbumin (Drewe, R. H. & Winzor, D. J.) 737–741
Phosphate, inorganic, reaction of, with the acyl-holoenzyme of pig skeletal-muscle D-glyceraldehyde 3-phosphate dehydrogenase (Armstrong, J. McD. & Trentham, D. R.) 513–527
Phosphatidate phosphohydrolase, role of, in the regulation of the biosynthesis of triacylglycerols in rat parametrial adipose-tissue homogenates (Christie, W. W., Hunter, M. L. & Vernon, R. G.) 571–577
Phosphatidylcholine, role of, in the binding of adenosine diphosphate to dogfish rectal-gland sodium-plus-potassium ion-dependent adenosine triphosphatase (Jensen, J. & Ottolenghi, P.) 815–817
Phosphodiesterase, alkaline, location of the activities of nucleotide pyrophosphatase and, on mouse lymphocyte plasma membrane (Abney, E. R., Evans, W. H. & Parkhouse, R. M. E.) 293–299
6-Phosphogluconate dehydrogenase, absorption procedure for the preparation of monospecific antisera for, and other specific enzymes for studies on enzyme turnover (Walker, J. H., Betts, S. A., Manning, R. & Mayer, R. J.) 355–362

Phospholipase A, actions of snake venom, trypsin and, *in vitro* on purified rat myelin (Banik, N. L., Gohil, K. & Davison, A. N.) 273–277
Phospholipid vesicles, modification of the activity of rabbit skeletal-muscle glyceraldehyde 3-phosphate dehydrogenase by adsorption on (Wooster, M. S. & Wrigglesworth, J. M.) 627–631
Pigeon, interaction of Cibacron Blue F3GA with liver nicotinamide–adenine dinucleotide kinase from (Apps, D. K. & Gleed, C. D.) 441–443
Pituitary gland, anterior, ox, effects of carboxymethylation of methionine residues in lutropin from, on the binding activity with receptor sites and interactions between subunits (Cheng, K.-W.) 71–77
Pituitary gland, anterior, ox, location of methionine residues in lutropin from, that are preferentially modified by carboxymethylation (Cheng, K.-W.) 79–87
Pituitary gland, anterior, ox, purification and properties of follitropin from (Cheng, K.-W.) 651–659
Placenta, human, full-term, purification and properties of arginase from (Porta, R., Esposito, C., Martin, A. & Della Pietra, G.) 579–583
Placenta, human, purification and chemical characterization of hexosaminidases A and B from (Lee, J. E. S. & Yoshida, A.) 535–539
Placenta, human, subunit structure of hexosaminidase A from (Beutler, E., Yoshida, A., Kuhl, W. & Lee, J. E. S.) 541–543
Placenta, rabbit, purification and characterization of lactogen from (Bolander, F. F., Jr. & Fellows, R. E.) 775–782
Plasma, cebus-monkey, comparison of the immunological and electrophoretic properties of 25-hydroxycholecalciferol-binding proteins of, and of plasma of various other monkeys and primates (Bouillon, R., Van Baelen, H. & De Moor, P.) 463–466
Plasma, human, properties of the primary inhibitor of plasmin in (Müllertz, S. & Clemmensen, I.) 545–553
Plasma membrane, *see* Membrane, plasma
Plasma, ox, isolation and characterization of an active three-chain species of thrombin from (Giglio, J. R., Rossi, A., Leone, F. A., Chiericato, G. & Say, J. C.) 29–33
Plasma, rat, isolation and physicochemical properties of acute-phase α_2-macroglobulin from (Gauthier, F. & Mouray, H.) 661–665
Plasma, rat, isolation and properties of acute-phase α_1- and α_2-macroglobulins from (Gordon, A. H.) 643–650
Plasmin, properties of the primary inhibitor of, in human plasma (Müllertz, S. & Clemmensen, I.) 545–553
Polyacrylamide gels, instantaneous monitoring of, for zones containing proteins during electrophoresis (Elliott, A.) 743–748
Polyadenylic acid, stability, ribonucleoprotein form and content of, of artichoke-tuber nuclear ribonucleic acid (Chapman, K. S. R. & Ingle, J.) 585–600
Polychaete worm (*Eunice aphroditois*), isolation, characterization and oxygen equilibrium of an extracellular haemoglobin from (Bannister, J. V., Bannister, W. H., Anastasi, A. & Wood, E. J.) 35–42
Polyguluronate lyase, purification and properties of, from a marine *Pseudomonas* sp. (Davidson, I. W., Sutherland, I. W. & Lawson, C. J.) 707–713

Polypeptide portions, lysozyme, chicken-egg, semi-synthesis of, by condensation of protected fragments (Rees, A. R. & Offord, R. E.) 487–493

Polypeptides, modified, preparation of protected fragments of chicken-egg lysozyme for the semisynthesis of (Rees, A. R. & Offord, R. E.) 467–479

Polypeptides, use of diazoalkanes as protecting agents in the semisynthesis of (Offord, R. E., Storey, H. T., Rees, A. R., Hayward, C. F., Johnson, W. H., Pheasey, M. H. & Wightman, D. A.) 480–486

Prolactin, purification and characterization of rabbit placental lactogen and its relationship to (Bolander, F. F., Jr. & Fellows, R. E.) 775–782

Protease, *see* Proteinase

Protein disulphide-isomerase, microsomal, liver, ox, differences between the activities of, and of glutathione–insulin transhydrogenase (Hawkins, H. C. & Freedman, R. B.) 385–393

Protein disulphide-isomerase, microsomal, liver, rat, comparison of the properties of, with those of glutathione–insulin transhydrogenase (Ibbetson, A. L. & Freedman, R. B.) 377–384

Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, purification and physical properties of the catalytic subunit of, from ox liver (Sugden, P. H., Holladay, L. A., Reimann, E. M. & Corbin, J. D.) 409–422

Protein kinases, adenosine 3′:5′-cyclic monophosphate-dependent, properties of, and other adenosine 3′:5′-cyclic monophosphate-binding proteins of various ox and rat tissues (Sugden, P. H. & Corbin, J. D.) 423–437

Protein–polysaccharides, *see* Proteoglycans

Proteinase inhibitor, α_1-, human, inactivation of human thrombin in the presence of (Matheson, N. R. & Travis, J.) 495–502

Proteinase inhibitor (α-1-antitrypsin), isolation and characterization of, from rhesus-monkey serum (Berninger, R. W. & Mathis, R. K.) 95–104

Proteins, accessibility of dinitrophenyl groups bound to, to univalent fragments of rabbit anti-dinitrophenyl antibody (Knight, C. G. & Green, N. M.) 323–333

Proteins, adenosine 3′:5′-cyclic monophosphate-binding, properties of, of various ox and rat tissues (Sugden, P. H. & Corbin, J. D.) 423–437

Proteins, alternative products in the reaction of 2-nitro-5-thiocyanatobenzoic acid with thiol groups of (Price, N. C.) 177–180

Proteins, electrophoretic method for the study of the reversible binding of ions to (Drewe, R. H. & Winzor, D. J.) 737–741

Proteins, 25-hydroxycholecalciferol-binding, plasma, comparison of the immunological and electrophoretic properties of, from cebus monkeys and from various other monkeys and primates (Bouillon, R., Van Baelen, H. & De Moor, P.) 463–466

Proteins, identification of, and their subunits of pig kidney-cortex microvillus membrane after sodium dodecyl sulphate/polyacrylamide-gel electrophoresis (Booth, A. G. & Kenny, A. J.) 395–407

Proteins, instantaneous monitoring of polyacrylamide gels for zones containing, during electrophoresis (Elliott, A.) 743–748

Proteins, loss of a homologous group of, from keratin from mice with the dominantly inherited Naked trait (Tenenhouse, H. S. & Gold, R. J. M.) 149–157

Proteins, method for calculating the relative contents of, in the major components of keratin from their amino acid compositions (Gold, R. J. M., Tenenhouse, H. S. & Adler, L. S.) 157–160

Proteoglycans, use of binding to, in a method for the determination of hyaluronate in the presence of other glycosaminoglycans and its application to human intervertebral disc (Hardingham, T. E. & Adams, P.) 143–147

Protons, formation constants of complexes of adenyl-5′-yl imidodiphosphate with bivalent metal ions and (Pettit, L. D. & Siddiqui, K. F.) 169–171

Pseudomonas sp., marine, purification and properties of alginate lyase from (Davidson, I. W., Sutherland, I. W. & Lawson, C. J.) 707–713

Pyruvate dehydrogenase complex, *Escherichia coli* K12, evidence for the presence of two lipoic acid residues per chain of lipoate acetyltransferase in (Danson, M. J. & Perham, R. N.) 677–681

Pyruvate kinase, skeletal-muscle, rabbit, identification of the active-site cysteine residue of, as the reactive group alkylated by 5-chloro-4-oxopentanoate (Chalkley, R. A. & Bloxham, D. P.) 213–219

Pyruvate kinase, skeletal-muscle, rabbit, selective modification of, by 5-chloro-4-oxopentanoate (Bloxham, D. P. & Chalkley, R. A.) 201–211

Rectal gland, dogfish, role of lipid in the binding of adenosine diphosphate to sodium-plus-potassium ion-dependent adenosine triphosphatase from (Jensen, J. & Ottolenghi, P.) 815–817

Reticulum, sarcoplasmic, skeletal-muscle, rabbit, binding of nucleotides and bivalent cations to calcium-plus-magnesium ion-dependent adenosine triphosphatase from (Yates, D. W. & Duance, V. C.) 719–728

all-*trans*-β-Retinoyl phosphate, chemical synthesis of (Frot-Coutaz, J. P. & De Luca, L. M.) 799–801

Rhesus monkey (*Macaca mulatta*), isolation and characterization of α-1-antitrypsin from serum from (Berninger, R. W. & Mathis, R. K.) 95–104

Ribonucleic acid, messenger, stability, polyadenylic acid content and ribonucleoprotein form of artichoke-tuber nuclear ribonucleic acid and its relationship to (Chapman, K. S. R. & Ingle, J.) 585–600

Ribonucleic acid, nuclear, artichoke-tuber, stability, polyadenylic acid content and ribonucleoprotein form of (Chapman, K. S. R. & Ingle, J.) 585–600

Ribonucleic acid, transfer, cysteine, baker's-yeast, application of a spectrophotometric method to the determination of the composition of oligonucleotides obtained by ribonuclease digestion of (Holness, N. J. & Atfield, G.) 15–22

Ribonucleic acids, messenger, model for the cytosol-governed regulation of the transcription of gene deoxyribonucleic acids in the biosynthesis of (Lichtenstein, A. V. & Shapot, V. S.) 783–794

Ribonucleic acids, transfer, *Escherichia coli* and rat liver, inhibitory effects *in vitro* of some amino acid analogues on the aminoacylation of (Old, J. M. & Jones, D. S.) 503–511

Ribonucleoprotein, stability, polyadenylic acid content and presence in the form of, of artichoke-tuber nuclear ribonucleic acid (Chapman, K. S. R. & Ingle, J.) 585–600
Ribosomal subparticles, large and small, construction and analysis of sucrose density gradients for use in the separation of, and components of other systems in zonal rotors (Hirst, W. & Cox, R. A.) 259–265
Ribulose 1,5-diphosphate carboxylase, plant, model for the kinetics of the activation of and of the carboxylation and oxygenation reactions catalysed by (Laing, W. A. & Christeller, J. T.) 563–570
Ribulose 1,5-diphosphate, model for the kinetics of the activation of plant ribulose 1,5-diphosphate carboxylase and of the carboxylation and oxygenation, of, catalysed by the enzyme (Laing, W. A. & Christeller, J. T.) 563–570

Saccharomyces cerevisiae, application of a spectrophotometric method to the determination of the composition of oligonucleotides obtained by ribonuclease digestion of cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 15–22
Saccharomyces cerevisiae, solubilization and other studies on adenylate cyclase from (Varimo, K. & Londesborough, J.) 363–370
Sarcoplasmic reticulum, *see* Reticulum, sarcoplasmic
Serine–pyruvate aminotransferase, identity of isoenzyme 1 of rat liver histidine–pyruvate aminotransferase with (Noguchi, T., Okuno, E. & Kido, R.) 607–613
Serum albumin, *see* Albumin, serum
Serum, cebus-monkey, comparison of the immunological and electrophoretic properties of 25-hydroxycholecalciferol-binding proteins of, and of serum from various other monkeys and primates (Bouillon, R., Van Baelen, H. & De Moor, P.) 463–466
Serum, rat, isolation and physicochemical properties of acute-phase α_2-macroglobulin from (Gauthier, F. & Mouray, H.) 661–665
Serum, rat, isolation and properties of acute-phase α_1- and α_2-macroglobulins from (Gordon, A. H.) 643–650
Serum, rhesus-monkey, isolation and characterization of α-1-antitrypsin from (Berninger, R. W. & Mathis, R. K.) 95–104
Sialic acid, formation of a stable chromophore of thiobarbituric acid in the presence of dimethyl sulphoxide and its application to the assay of, in analytical de-*O*-acetylation of glycoproteins (Skoza, L. & Mohos, S.) 457–462
Sialoproteins, formation of a stable chromophore of thiobarbituric acid in the presence of dimethyl sulphoxide and its application to the assay of sialic acid in analytical de-*O*-acetylation of (Skoza, L. & Mohos, S.) 457–462
Skeletal muscle, *see* Muscle, skeletal
Small intestine, *see* Intestine, small
Snake (*Naja naja*) venom, actions of phospholipase A, trypsin and, *in vitro* on purified rat myelin (Banik, N. L., Gohil, K. & Davidson, A. N.) 273–277
Somatotropin, *see* Growth hormone
Sperm whale, kinetics and mechanism of the recombination reaction between apomyoglobin from, and haemin (Adams, P. A.) 371–376
Steroids, $\Delta^{8(14)}$-, identification of, in *Methylococcus capsulatus* (Bouvier, P., Rohmer, M., Benveniste, P. & Ourisson, G.) 267–271
Sterols, $\Delta^{8(14)}$-, identification of, in *Methylococcus capsulatus* (Bouvier, P., Rohmer, M., Benveniste, P. & Ourisson, G.) 267–271
Streptomyces fradiae, role of the pseudo-disaccharide neamine as an intermediate in the biosynthesis of neomycin in *Streptomyces rimosus* forma *paromomycinus* and (Pearce, C. J., Barnett, J. E. G., Anthony, C., Akhtar, M. & Gero, S. D.) 601–606
Streptomyces rimosus forma *paromomycinus*, role of the pseudo-disaccharide neamine as an intermediate in the biosynthesis of neomycin in *Streptomyces fradiae* and (Pearce, C. J., Barnett, J. E. G., Anthony, C., Akhtar, M. & Gero, S. D.) 601–606
Subcomponent C1q, complement, human, circular-dichroism and electron-microscopy studies before and after limited proteolysis by pepsin of (Brodsky-Doyle, B., Leonard, K. R. & Reid, K. B. M.) 279–286
Submitochondrial particles, heart, ox, reaction of 4-chloro-7-nitrobenzofurazan with the adenosine triphosphatase of (Ferguson, S. J., Lloyd, W. J. & Radda, G. K.) 347–353
Sucrose, construction and analysis of density gradients of, for use in zonal rotors (Hirst, W. & Cox, R. A.) 259–265
Sugar sulphate esters, substrate specificity of *Charonia lampas* arylsulphatase and glycosulphatase towards (Hatanaka, H., Ogawa, Y. & Egami, F.) 445–448
Sulphydryl, *see* Thiol
Supernatant fraction, *see* Cytosol
(±)-Synephrine, conversion of, into *p*-hydroxyphenylacetaldehyde catalysed by an enzyme from *Arthrobacter synephrinum* (Veeraswamy, M., Abitha Devi, N., Krishnan Kutty, R. & Subba Rao, P. V.) 807–809

Tendon, ox, comparison of the compositions of, and bone matrix with particular reference to the content of glycoproteins (Herring, G. M.) 749–755
Testis, rabbit, purification and properties of arylsulphatase A from (Yang, C.-H. & Srivastava, P. N.) 133–142
Tetrahymena pyriformis (Phenoset A), amino acid sequence of cytochrome *c* from (Tarr, G. E. & Fitch, W. M.) 193–199
Thiobarbituric acid, formation of a stable chromophore of, in the presence of dimethyl sulphoxide and its application to the assay of sialic acid in analytical de-*O*-acetylation of glycoproteins (Skoza, L. & Mohos, S.) 457–462
Thiol group, active-site, identification of, of rabbit skeletal-muscle pyruvate kinase as the reactive group alkylated by 5-chloro-4-oxopentanoate (Chalkley, R. A. & Bloxham, D. P.) 213–219
Thiol groups, alternative products in the reaction of 2-nitro-5-thiocyanatobenzoic acid with, of proteins (Price, N. C.) 177–180
Thiol groups, use of 2,2′-dipyridyl disulphide as a thiol titrant and reactivity probe in an investigation of, of high-activity urease prepared from jack-bean meal (Norris, R. & Brocklehurst, K.) 245–257
Thiol–protein disulphide oxidoreductases, microsomal, liver, ox, differences between the activities of (Hawkins, H. C. & Freedman, R. B.) 385–393

Thiol–protein disulphide oxidoreductases, microsomal, liver, rat, activities and comparison of the properties of (Ibbetson, A. L. & Freedman, R. B.) 377–384
Thiols, reactions of D-glyceraldehyde 3-phosphate with, and with pig skeletal-muscle D-glyceraldehyde 3-phosphate dehydrogenase (Armstrong, J. McD. & Trentham, D. R.) 513–527
Thrombin, human, inactivation of, in the presence of α_1-proteinase inhibitor (Matheson, N. R. & Travis, J.) 495–502
Thrombin, isolation and characterization of an active three-chain species of, from ox plasma (Giglio, J. R., Rossi, A., Leone, F. A., Chiericato, G. & Say, J. C.) 29–33
Transcalciferin, *see* 25-Hydroxycholecalciferol-binding protein
Transcription, model for the cytosol-governed regulation of, of gene deoxyribonucleic acids in the biosynthesis of messenger ribonucleic acids (Lichtenstein, A. V. & Shapot, V. S.) 783–794
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Triacylglycerols, biosynthesis of, in rat parametrial adipose-tissue homogenates (Christie, W. W., Hunter, M. L. & Vernon, R. G.) 571–577
Triglycerides, *see* Triacylglycerols
Troponin I, heart-muscle, rabbit, amino acid sequence of (Grand, R. J. A., Wilkinson, J. M. & Mole, L. E.) 633–641
Trypsin, actions of snake venom, phospholipase A and, *in vitro* on purified rat myelin (Banik, N. L., Gohil, K. & Davison, A. N.) 273–277
Trypsin inhibitor (α-1-antitrypsin), isolation and characterization of, from rhesus-monkey serum (Berninger, R. W. & Mathis, R. K.) 95–104
Trypsin inhibitor, α_1-, human, inactivation of human thrombin in the presence of (Matheson, N. R. & Travis, J.) 495–502
Tryptophan residues, determination of the number and relative positions of, in serum albumins from various animal species (Feldhoff, R. C. & Peters, T., Jr.) 529–533
Tubers, artichoke, stability, polyadenylic acid content and ribonucleoprotein form of nuclear ribonucleic acid from (Chapman, K. S. R. & Ingle, J.) 585–600

Urease, preparation by covalent chromatography of, of high activity from jack-bean meal and use of 2,2′-dipyridyl disulphide as a thiol titrant and reactivity probe in an investigation of its thiol groups (Norris, R. & Brocklehurst, K.) 245–257
Uridine diphosphate glucose pyrophosphorylase, liver, ox, binding of ligands to (Stevens, R. A. J. & Phelps, C. F.) 65–70
Uridine 5′-methylenediphosphoro-P^2-phosphate, synthesis of, and its binding to ox liver uridine diphosphate glucose pyrophosphorylase (Stevens, R. A. J. & Phelps, C. F.) 65–70

Venom, snake (*Naja naja*), actions of phospholipase A, trypsin and, *in vitro* on purified rat myelin (Banik, N. L., Gohil, K. & Davison, A. N.) 273–277
Vitamin D_3, *see* Cholecalciferol

Worm, polychaete (*Eunice aphroditois*), isolation, characterization and oxygen equilibrium of an extracellular haemoglobin from (Bannister, J. V., Bannister, W. H., Anastasi, A. & Wood, E. J.) 35–42

Yeast, baker's (*Saccharomyces cerevisiae*), application of a spectrophotometric method to the determination of the composition of oligonucleotides obtained by ribonuclease digestion of cysteine transfer ribonucleic acid from (Holness, N. J. & Atfield, G.) 15–22
Yeast, baker's (*Saccharomyces cerevisiae*), solubilization and other studies on adenylate cyclase from (Varimo, K. & Londesborough, J.) 363–370
Yeast (*Candida utilis* C.E.C.T. 1061), purification and properties of exo-β-D-glucanase from cell-free extracts of (Notario, V., Villa, T. G. & Villanueva, J. R.) 555–562

Zonal rotors, construction and analysis of sucrose density gradients for use in (Hirst, W. & Cox, R. A.) 259–265